

## FAQ

9 April 2024

### **Selected questions and answers relating to hygiene of food and consumables in times of the bird flu – How can I protect myself and my family?**

→ Changes to the version dated 16 March 2022: Addition of information on the transmission of the bird flu virus from infected cattle to humans through contact with the animals and consumption of milk

---

Concerned persons are asking whether poultry, poultry products or other foods can become contaminated with the bird flu virus (various subtypes of the avian influenza virus including H5N1 and H5N8). Here, the German Federal Institute for Risk Assessment (BfR) provides answers to these questions.

Generally, the transmission of the pathogen via infected food cannot be ruled out. However, infections of humans with the bird flu virus are rare, and direct and close contact with infected birds or other animal species appears to be the main transmission route to humans. The BfR does not have any data verifying that humans have been infected and became ill after ingestion of food that was contaminated with the bird flu virus.

As the virus is sensitive to high temperatures, no adverse health effects are to be expected if the food has been well cooked. Meat is considered well cooked, if all sides and the core reach a temperature of at least 70 °C for 2 minutes. This is evident when the poultry meat is no longer red or pink and no red juices are leaking out.

So far, there is no evidence suggesting that humans can be infected by raw eggs or raw sausage products containing poultry meat from infected animals. However, it has been proven that eggs of infected animals can contain the virus both on the shell and also in the egg white and yolk. As a precaution, anyone who wants to protect themselves against bird flu viruses and other pathogens that are possibly present in eggs and egg products, should not consume raw egg products (whipped egg whites, tiramisu, etc.). The egg white and egg yolk of boiled eggs should be solid.

The consumption of pasteurised milk from cattle possibly infected with the bird flu virus is not expected to have any adverse health effects, as the virus - like other pathogenic microorganisms - is efficiently inactivated by heating.

### **What is bird flu?**

The highly pathogenic avian influenza (bird flu, classical fowl plaque) is a disease that is highly infectious for domestic fowl, especially chickens and turkeys that suffer from serious disease courses. Bird flu is caused by various avian flu strains, including the subtypes H5N1, H5N6, H5N8 and H7N9, that can also cause serious illness among humans in some cases. The BfR provided more information about the subtype H5N1 in 2004 at

[https://www.bfr.bund.de/cm/343/gefluegelpest\\_infektionsrisiko\\_fuer\\_den\\_verbraucher\\_durch\\_lebensmittel.pdf](https://www.bfr.bund.de/cm/343/gefluegelpest_infektionsrisiko_fuer_den_verbraucher_durch_lebensmittel.pdf)

and about subtype H5N8 in 2021 at

<http://vm-webextern-m.bfr.bund.de/cm/349/current-avian-influenza-case-in-germany-virus-transmission-h5n8-from-poultry-food-consumption-is-unlikely.pdf>

Further information is available on the websites of the Friedrich Loeffler Institute ([www.fli.de](http://www.fli.de)), the Robert Koch Institute ([www.rki.de](http://www.rki.de)), and the Federal Ministry of Food and Agriculture ([www.bmel.bund.de](http://www.bmel.bund.de)).

### **Where has the bird flu been detected so far?**

There is more information about the latest spread of bird flu on the homepage of the Food and Agriculture Organization of the United Nations (FAO) at <https://www.fao.org/animal-health/situation-updates/en> and the World Health Organization (WHO), at [https://www.who.int/health-topics/influenza-avian-and-other-zoonotic#tab=tab\\_1](https://www.who.int/health-topics/influenza-avian-and-other-zoonotic#tab=tab_1).

### **Can the bird flu virus be transmitted to humans?**

It is difficult to transmit the bird flu virus to humans. The WHO received a total of 882 reports of human infection with avian influenza viruses of subtype H5N1 from 18 countries between January 2003 and December 2023 (WHO, Avian Influenza Weekly Update Number 938 dated 15.03.2024). This is a very low number in light of the huge spread and high infection density of the pathogen in the global poultry population and the associated frequent contacts between humans and poultry. Also, only few cases of human infection with the other subtypes have been reported (WHO, Avian Influenza Weekly Update Number 938 dated 15.03.2024).

### **How can the bird flu virus be transmitted to humans?**

The bird flu virus is primarily transmitted by breathing in (inhaling) contaminated dust particles or droplets. The virus can possibly also be transmitted by smear infection with virus-contaminated excretions on mucous membranes. Persons who have close contact with

poultry or diseased mammals should take suitable precautions. Detailed recommendations about this are stated on the websites of the Federal Institute for Occupational Safety and Health (<http://www.baua.de>) and the Robert Koch Institute (<http://www.rki.de>).

### **Can the bird flu virus be transmitted to humans through infected cattle and their milk?**

In March 2024, cases of H5N1 infection in cattle were reported in several US states. In this context, an H5N1 infection was also detected in a human with close contact to the diseased cattle. This was a strain of the virus that is currently widespread worldwide, particularly in wild birds. In the USA, the virus was detected in milk samples and swab samples from the mouth and throat of sick cattle. The disease in cattle was mainly characterised by temporary fever, lethargy, reluctance to eat and a drop in milk yield. Transmission of the virus between cattle cannot be ruled out at present. The infected human patient showed mild symptoms of the disease, mainly conjunctivitis. So far there is no further evidence of human-to-human transmission. Genome analyses of the virus strain also revealed no evidence of changes that would suggest an increased risk to humans.

According to the reports, the virus was also detected in milk samples from sick cattle. It is not yet known whether humans can become infected by consuming H5N1-contaminated milk. However, previous human H5N1 infections were mostly due to close contact with sick animals. The consumption of pasteurised milk from cattle possibly infected with the bird flu virus is not expected to have any adverse health effects, as the virus - like other pathogenic micro-organisms - is efficiently inactivated by heating. Nevertheless, milk from infected animals should not be sold. Raw milk should generally be heated before consumption, also to protect against other pathogens that may be present.

### **Is there a risk of infection when preparing and consuming food?**

Based on the current knowledge, direct contact with poultry is the most important transmission route of the virus between poultry and humans. There is only little known about virus transmission by the consumption of raw poultry meat products from infected animals. However, with respect to the preventive consumer protection, care should be taken to comply with hygiene rules when handling or preparing raw poultry meat and poultry meat products. As the virus is highly sensitive to high temperatures, no adverse health effects are to be expected if the food has been well cooked. Meat is considered well cooked, if all sides and the core reach a temperature of at least 70 °C for 2 minutes. This is evident to consumers when the poultry meat is no longer red or pink and no red juices are leaking out. In the case of eggs or egg products, it is possible to protect against these viruses and other possibly present pathogens by avoiding the consumption of raw egg products (whipped egg whites, tiramisu etc.), and by ensuring that the egg white and yolks of boiled eggs are solid. The consumption of pasteurised milk from cattle possibly infected with the bird flu virus is not expected to have any adverse health effects, as the virus - like other pathogenic microorganisms - is efficiently inactivated by heating. Raw milk should generally be heated before consumption, also to protect against other pathogens that may be present.

### **How can consumers protect themselves?**

Humans should avoid contact with wild birds. This particularly applies for died animals and for regions in which the bird flu virus was detected in dead wild birds (restricted areas, observation areas). Visitors to countries in which the bird flu is widely distributed among poultry flocks should avoid direct contact with poultry and their excretions. It is advised to refrain from visiting poultry markets or poultry farms. Import bans on poultry or other birds, poultry meat, eggs and other poultry products, or feathers or untreated hunting trophies from affected countries into the European Union should always be observed to prevent further outbreaks of bird flu in domestic commercial livestock. When preparing poultry meat and raw eggs, the general hygiene regulations should be observed:

- Store and prepare raw poultry products and other food separately, especially if the latter is not reheated.
- Thoroughly clean equipment and surfaces that have come into contact with raw poultry products with warm water and detergent.
- Dispose of packaging materials, thawing water and similar immediately.
- Wash hands with warm water and soap.
- Cook poultry meals thoroughly. This means that all areas, even the core, must have reached a temperature of at least 70 °C for 2 minutes.
- In the case of eggs or egg products, it is possible to protect against these viruses and other possibly present pathogens by avoiding the consumption of raw egg products (whipped egg whites, tiramisu etc.), and by ensuring that the egg white and yolks of boiled eggs are solid.

In countries where the virus has also been detected in cattle, close contact with infected animals should be avoided. Raw milk should generally be heated before consumption, also to protect against other pathogens that may be present.

### **Can animals and humans become infected with bird flu from animal or human drinking water?**

Surface water from waters with large populations of wild birds may, in principle, be contaminated with bird flu viruses. It should not be used as drinking water for breeding and commercial poultry, if it has not been appropriately treated.

Drinking water in Germany is primarily collected from deep wells. This guarantees a high level of safety regarding microbiological risks. Wherever drinking water is collected from surface waters, complex technical procedures are used that comply with the tolerances and requirements of the Drinking Water Regulations. This reliably rules out contamination of drinking water with bird flu viruses. The German Federal Office for Environment provides more information about drinking water

(<https://www.umweltbundesamt.de/themen/wasser/trinkwasser>).

### **Can lettuce or vegetables transmit bird flu viruses?**

In principle, lettuce, vegetable and fruit farmed outdoors can become contaminated with bird droppings. These may contain bird flu viruses. To minimise the risk of transmission of viruses, bacteria and parasites to humans, lettuce, fruit and vegetables are roughly cleaned in the production facility before storage. The usual hygiene rules should be observed when preparing lettuce, fruit and vegetables at home: Before preparation or consumption, wash thoroughly to remove all earth and manure particles. Cooked vegetables also offer more safety against infection: The bird flu virus is inactivated, if heated to above 70 °C for at least 2 minutes.

### **How can consumers recognise the origin of eggs or poultry?**

The code number of the packing point must be stated on egg cartons. This may also contain information about the origin of the eggs. The eggs themselves bear the country code of the country of origin.

Poultry meat sold in retail shops bears the approval number of the slaughterhouse or meat processing facility, and poultry meat that was imported from non-EU countries the country of origin, on the packaging or label.

There are relevant legal regulations that govern the import and the trade of live poultry from livestock populations and their products. There are more recommendations about hygienic handling of poultry meat on the websites of the German Federal Institute for Risk Assessment ([www.bfr.bund.de](http://www.bfr.bund.de)).

### **Can humans become infected with the bird flu virus after eating shellfish and fish?**

It is known that viruses can be transmitted to humans after eating uncooked or inadequately cooked shellfish and thereafter cause gastroenteritis. These are usually norovirus and hepatitis A virus that are very resistant to environmental influences. In contrast, influenza viruses are very sensitive to environmental influences. Examinations of samples from surface waters confirm the tendency that the infectivity of avian influenza viruses decreases proportionally to the salt content. Human infection with bird flu viruses after eating shellfish or fish is therefore unlikely and has not yet been proven. Cooking shellfish and fish offers additional protection against infection: The bird flu virus is usually inactivated, if heated to above 70 °C for at least 2 minutes.

### **Can poultry manure be still used as fertilizer?**

Poultry manure from healthy poultry livestock not infected with the bird flu virus from operations that lie outside the bird flu restricted areas can still be used. Poultry manure made of infected or potentially infected stocks must be 'decontaminated', i.e. disinfected. There are a series of effective procedures that are specified by the Tiergesundheitsgesetz and related regulations.

**Can humans become infected with bird flu from objects like down jackets, pillows or quilts produced with feathers or down from infected animals?**

Down is washed during production and then dried at 100 °C. Influenza viruses are very sensitive to heat and to the detergents used in soaps and washing agents. Based on the current knowledge, the methods used to produce down also inactivate bird flu viruses so that they are no longer infectious. An infection with the bird flu virus through objects that contain feathers or down is therefore very unlikely.

## About the BfR

The German Federal Institute for Risk Assessment (BfR) is a scientifically independent institution within the portfolio of the Federal Ministry of Food and Agriculture (BMEL) in Germany. The BfR advises the Federal Government and the States ('Laender') on questions of food, chemicals and product safety. The BfR conducts independent research on topics that are closely linked to its assessment tasks.

*This text version is a translation of the original German text which is the only legally binding version.*

### Legal notice

Publisher:

**German Federal Institute for Risk Assessment**

Max-Dohrn-Straße 8-10

10589 Berlin, Germany

T +49 30 18412-0

F +49 30 18412-99099

bfr@bfr.bund.de

bfr.bund.de/en

Institution under public law

Represented by the president Professor Dr Dr Dr h.c. Andreas Hensel

Supervisory Authority: Federal Ministry of Food and Agriculture

VAT ID No. DE 165 893 448

Responsible according to the German Press Law: Dr Suzan Fiack



valid for texts produced by the BfR

images/photos/graphics are excluded unless otherwise indicated)

**BfR** | Identifying Risks –  
Protecting Health